



- For highly sensitive drugs, leachables of the primary packaging into the drug product need to be minimized in order to protect their stability.
- The adsorption of proteins to the glass surface can result in conformational changes, denaturation, aggregation, and loss of effectiveness.
- SCHOTT Type I plus® vials have an ion barrier coating that can reduce adsorption and ensures low leachable levels for pharmaceutical products in the neutral and acidic pH ranges.
- Inner coating is applied using patented and proven plasma impulse chemical vapor deposition (PICVD) technology.
- An additional "release criterion" has been specifically developed.



SiO<sub>2</sub> layer applied via PICVD



Layer thickness of ~ 100–200 nm



Long-term layer stability



Coating bonds covalently to the glass substrate



Suitable for depyrogenation



Dense coating (i.e. non-porous)



Surface shows excellent barrier properties in reducing ion leaching: Na, Ca, B, Si, Al



Stable after washing process



Compliant with all current standards, such as Ph. Eur, USP, JP, and CP

#### Secured drug stability

Ion barrier provides stability for leachable sensitive drugs (e.g. radiopharmaceuticals and aluminum-sensitive drugs)

#### Reduced adsorption

For certain proteins, adsorption can be reduced thanks to a specialized inner surface\*

Avoidance of pH-shift
Particularly relevant to WFI




<sup>\*</sup> adsorption behavior needs to be tested case by case



#### 1) Leaching


Autoclaving filled with 0.1 M HCl (pH 1 at start).



# 2) Determination of sodium via AAS

Certified release criterion for sodium (Na) – defined limit value.

## Verifications: Avoidance of pH shift




#### Method

10 ml vials, washed in accordance with ISO 4802. Autoclaving 1 h at 121 °C. Storage at room temperature. Measurement of pH value prior to and after autoclaving (directly after, one day, one week, one/six/12/18 months).


#### Result

After autoclaving, standard Type I vials show pH shifts from 5.5 up to 7.5. SCHOTT Type I plus® vials fully prevent pH shift.

## Verifications: Secured drug stability



#### **Verifications: Reduced adsorption**



## Method

10R vials, filled with purified water, 1 h autoclaving at 121 °C. Leached ions in  $\mu g/ml$  by ICP-MS (ICP-OES for Si).

#### Result

The ion barrier is effective for all other elements of the glass matrix.

## Method

Nicotinic acetylcholine receptor, concentration: 150  $\mu$ g/ml, see: "A. Schrattenholz: Drug loss through adsorption of a nicotinic acetylcholine receptor in SCHOTT Type I plus® and standard vials".

# Result

Tested protein shows less tendency to adsorb to SCHOTT Type I plus® vials.

General ordering information												
Quality level	TopL	TopLine with additional release test										
Packaging		<ul> <li>Tray with optional divider</li> <li>Pre-washed and pre-sterilized: adaptiQ® (tray, cup nest)</li> </ul>										
Palletizing		Standard Euro pallet (1200 x 800 mm) contains $15-27$ layers of nine trays each										
Formats	2R	3R	4R	6R	8R	10R	15R	20R	25R	30R	50R	
Pieces per tray	344	344	344	186	186	154	154	95	95	95	40	

Many configurations are available in small quantities as "Fast Track Articles".

Visit our online shop or speak to your sales representative for more information.

schott-pharma.com/vials

SCHOTT Pharma AG & Co. KGaA, Hattenbergstrasse 10, 55122 Mainz, Germany pharma@schott.com



ENGLISH 07/2024 kn/nino Printed in Germany